



Category Theory


Roy L. Grole

University of Leicester. 2018.

M&S

Examples of Categories





e.g. $F(x_0) \circ K \stackrel{\text{def}}{=} F(x_0)[K/x_0]$

$$t := x_0 \mid K \mid F(t) \mid G(t, t)$$

WHAT IS A CATEGORY?

- A theory of "typed functions"
- "typed functions"
 - * possess identities
 - * are composable
 - * compose associatively

-
- remark: objects are "recoverable" from the identities
 - regard the above very informally!

General Definition of Category $\mathcal{C}$

- See slide for sets...
- Collection of **objects** $A, B, C \dots$

-

$$\mathcal{C}(A, B) =$$

$$\{\underline{f} \mid A \xrightarrow{f} B\}$$

collection of **morphisms** for each A, B

- satisfying **axioms** that define a category

More Examples

— Opposite of

is

$$0 \rightarrow 1 \rightarrow 2 \rightarrow 3 \rightarrow \dots$$

$$0 \leftarrow 1 \leftarrow 2 \leftarrow 3 \leftarrow \dots$$

identities,
transitivity
omitted

[isomorphic to $\dots \rightarrow -2 \rightarrow -1 \rightarrow 0$]

— $(X, \leq)$ "is" a category, say $\mathbb{C}$.


$$* \text{ ob } \mathbb{C} = X$$

$$* \mathbb{C}(x, x') = \begin{cases} \{*\} & \Leftrightarrow x \leq x' \\ \emptyset & \Leftrightarrow \text{otherwise} \end{cases}$$

— $\mathbb{C}^{\text{op}} = (X, \leq^{\text{op}})$ where

$$\forall x, x' \in X, x \leq^{\text{op}} x' \Leftrightarrow x' \leq x$$


A **monoid** is a category with one object $*$
 for example $(\text{list}(A), ++, [])$



$l, l' \in \text{list}(A)$ — $++$ append **associative** :
 — $[] ++ l = l = l ++ []$:

morphisms

... and Mon is a category (!)



— homomorphisms
compose



— identity function
is a homomorphism

WHAT IS A FUNCTOR?


- A mapping between categories that preserves the (fundamental) structure.
- maps objects to objects
- maps morphisms to morphisms
- preserves identities & composition

Examples of Functors

– $F : \underline{\text{Set}} \rightarrow \underline{\text{Mon}}$



– $F(\text{id}_A) = \text{map}(\text{id}_A) = \text{id}_{\text{list}(A)}$



$$\begin{aligned}
 [a_1, a_2] &\mapsto \text{map}(g \circ f) [a_1, a_2] \\
 &= [(g \circ f) a_1, (g \circ f) a_2] \\
 &= [g(f(a_1)), g(f(a_2))] \\
 &= \text{map}(g) \circ \text{map}(f) [a_1, a_2]
 \end{aligned}$$

$\oplus \checkmark$

General Definition of Functor $F: \mathbb{C} \rightarrow \mathbb{D}$


$$- \quad A \quad \vdash \quad F A$$

– $G_A \xrightarrow{id_A} G_{FA}$

The diagram shows two commutative triangles. The left triangle has vertices A , B , and C . A curved arrow f goes from A to B , a curved arrow g goes from B to C , and a dashed arrow $g \circ f$ goes from A to C . A pink circle with a cross is in the center. An arrow points to the right triangle, which has vertices FA , FB , and FC . A curved arrow Ff goes from FA to FB , a curved arrow Fg goes from FB to FC , and a dashed arrow $F(g \circ f)$ goes from FA to FC . A pink circle with a cross is in the center. Below the right triangle, the equation $F(g \circ f) = (Fg) \circ (Ff)$ is written.

Example of Natural Transformation

- $\text{rev} : F \rightarrow F$ ← - - - Think: polymorphic function 😊
- specified by $\text{rev}_A : FA \rightarrow FA$
- such that





$$Ff \stackrel{\text{def}}{=} \text{map}(f)$$

$$FA \stackrel{\text{def}}{=} \text{list}(A)$$

$$(id_F)_A \stackrel{\text{def}}{=} id_{FA}$$


$$(\beta \circ \alpha)_A \stackrel{\text{def}}{=}$$



C — objects $F, G, H \dots$
 D — morphisms $\alpha, \beta, \gamma, \delta \dots$

SUMMARY

CATEGORIES, FUNCTORS, NATURAL TRANSFORMATIONS



- $\circ$ associative
- $\circ$ id unitary

Universal Properties : Structure up to Isomorphism

$$\underline{\forall X. \exists!} \quad \begin{array}{l} f_X : X \longrightarrow T \\ g_X : X \longrightarrow T' \end{array}$$

Hence



$$\begin{array}{ccccc} T' & \xrightarrow{\exists f_{T'}} & T & \xrightarrow{\exists g_T} & T' \\ & \searrow & \downarrow \text{red} & \nearrow & \\ & & g_T \circ f_{T'} & & \\ & \text{red} \parallel & & & \\ & \text{red} ! & & & \\ & \text{red} id_{T'} & & & \end{array}$$

& via a symmetrical argument

$$T \cong T'$$

Example of Binary Product of A, B


$$A = \{2, 4\} \quad B = \{5, 6\} \quad C = \{\alpha, \beta\}$$



$\langle \hat{2}, \hat{5} \rangle$
is the
unique
morphism
such that

$$\langle \hat{2}, \hat{5} \rangle(x) = (\hat{2}(x), \hat{5}(x)) = (2, 5)$$


Example of Binary Coproduct of A, B




$$[\hat{\alpha}, \hat{\beta}](\xi) = \text{case } \xi \text{ of } \begin{array}{l} 2, 4 \mapsto \alpha \\ 5, 6 \mapsto \beta \end{array}$$

Example of Binary Coproduct in PreSet

$$X + Y = \{(x, 1), (y, 2) \mid x \in X, y \in Y\}$$



Illustrating Adjunctions $L \dashv R$



a natural bijection

$\mathbb{C}(-, R+), \mathbb{D}(L^{\text{op}}-, +) : \mathbb{C}^{\text{op}} \times \mathbb{D} \rightarrow \underline{\text{Set}}$
are isomorphic in $\underline{\text{Set}}^{\mathbb{C}^{\text{op}} \times \mathbb{D}}$:

— there are natural transformations
 $\Phi_{(A,B)} : \mathbb{C}(A, R B) \xrightleftharpoons{\quad} \mathbb{D}(L^{\text{op}} A, B) : \Psi_{(A,B)}$
 such that $\cong$


$$\begin{array}{ccccccc}
 LA' & \xrightarrow{L\phi} & LA & \xrightarrow{\bar{f}} & B & \xrightarrow{\psi} & B' \quad \text{in } \mathbb{D} \\
 A' & \xrightarrow{\phi} & A & \xrightarrow{f} & RB & \xrightarrow{R\psi} & RB' \quad \text{in } \mathbb{C}
 \end{array}$$

$$\begin{array}{ccc}
 \mathbb{C}(A, RB) & \xrightarrow{\quad} & \mathbb{D}(LA, B) \\
 \downarrow \mathbb{C}(\phi, R\psi) & \begin{array}{c} f \dashv \xrightarrow{\quad} \bar{f} \dashv \\ \downarrow \quad \quad \downarrow \\ R\psi \circ f \circ \phi \dashv \xrightarrow{\quad} R\psi \circ f \circ g \end{array} & \begin{array}{c} \bar{f} \dashv \xrightarrow{\quad} \psi \circ \bar{f} \circ L\phi \\ \downarrow \quad \quad \parallel \\ \mathbb{C}(L\phi, \psi) \end{array} \\
 \mathbb{C}(A', RB') & \xrightarrow{\quad} & \mathbb{D}(LA', B')
 \end{array}$$

Examples of Adjunctions

$list \rightarrow \mathcal{U}$

– $\mathcal{U} : \underline{Mon} \rightleftharpoons \underline{Set} : list$




$$f([a_1, a_2]) = f([a_1] ++ [a_2]) = f([a_1]) f([a_2]) = \bar{f}(a_1) \bar{f}(a_2)$$

$$\circ \circ f([a_1, a_2]) = \hat{f}([a_1, a_2]) \Rightarrow \cong_{bij}$$

$$- \quad \Delta : \underline{\text{Set}} \rightleftharpoons \underline{\text{Set}} \times \underline{\text{Set}} : \sqcap$$

$$\Delta \dashv \sqcap$$



$$(f, g)(x, y) = (fx, gy) = (\pi_A \circ \langle f, g \rangle x, \pi_B \circ \langle f, g \rangle y)$$

$$= (\pi_A \circ \overline{(f, g)} x, \pi_B \circ \overline{(f, g)} y) \circ (f, g)(x, y) = \widehat{(f, g)}(x, y) \Rightarrow \cong_{\text{bij}}$$

Example of Algebra and Homomorphism

- Take a monoid (M, b, e)
- $\hat{e} : 1 \rightarrow M$ $b : M \times M \rightarrow M$

Define^{*} $F(-) = 1 + (- \times -)$ and note

$$\begin{array}{ccc} 1 + (M \times M) & \xrightarrow{[\hat{e}, b]} & M \\ * \quad \downarrow \hat{e} & & \downarrow e \\ & \xrightarrow{b} & \end{array}$$

$$(m, n) \mapsto m \cdot n$$

$$FM \xrightarrow{[\hat{e}, b]} M \text{ is an algebra}$$

* what is the effect on morphisms?

Example of Algebra and Homomorphism

— Similarly (S, β, ε) . Then $\theta : M \rightarrow S$ is a homomorphism $\Leftrightarrow \theta$ is a homomorphism of algebras

— homomorphism means

$$* \theta(e) = \varepsilon$$

$$* \theta(m \cdot n) = \theta(m) \cdot \theta(n)$$

$\Rightarrow$

FM

$$\boxed{1 + (M \times M)} \xrightarrow{[\hat{e}, b]} M$$

F θ
||

$1 + (\theta \times \theta)$

*

|

(m, n)

|

$m \circ n$

|

|

$\theta(m \circ n)$
||

$(\theta m, \theta n)$

|

$\theta m \circ \theta n$

✓

*

|

1

+

$(S \times S)$

$[\hat{e}, \beta]$

|

S


FS

e

e

θ

Natural Numbers as an Initial Algebra



CASE STUDY

The Mini Yoneda Lemma
for Type Theorists

– for details see slides...

plus...

$$\begin{array}{ccc}
 \mathbb{C}(A, A) & \xrightarrow{\rho_A} & \mathbb{C}(A, B) \\
 \downarrow \mathbb{C}(m, id_A) & \begin{array}{c} id_A \mapsto \rho_A(id_A) \\ \downarrow \\ id_A \circ m \mapsto \rho_A(id_A) \circ m \end{array} & \downarrow \mathbb{C}(m, id_B) \\
 & \xrightarrow{\rho_G} & \\
 \mathbb{C}(G, A) & \xrightarrow{\rho_G} & \mathbb{C}(G, B) \\
 \downarrow \mathbb{C}(n, id_A) & \begin{array}{c} m \mapsto \rho_G(m) \\ \downarrow \\ m \circ n \mapsto \rho_G(m) \circ n \end{array} & \downarrow \mathbb{C}(n, id_B) \\
 & \xrightarrow{\rho_{G'}} & \\
 \mathbb{C}(G', A) & \xrightarrow{\rho_{G'}} & \mathbb{C}(G', B)
 \end{array}$$

$$\circ \circ (4) \iff \rho_- : \mathbb{C}(-, A) \rightarrow \mathbb{C}(-, B) \text{ nat. trans.}$$

$$\iff \rho_{G'}(m_A) = \rho_A(id_A) \circ m_A \quad (\text{Why does } \Leftarrow \text{ hold?})$$

CASE STUDY

A ccc can be defined in terms of an adjunction

– for details see slides...

plus...

Cartesian Closed Categories

- For any B, C there is $B \Rightarrow C$ and $ev : (B \Rightarrow C) \times B \longrightarrow C$
- For any $f : A \times B \rightarrow C$ there is $\lambda f : A \rightarrow (B \Rightarrow C)$ *unique* such that

$$\begin{array}{ccc} A \times B & \xrightarrow{f} & C \\ \lambda f \times id_B \downarrow & \oplus & \nearrow ev \\ (B \Rightarrow C) \times B & & \end{array}$$

λf is the
currying of f

Example of Set

$$- \text{ev} : (B \Rightarrow C) \times B \longrightarrow C$$

$$(f, b) \longmapsto f(b)$$

$$\text{where } B \Rightarrow C = \{f \mid f: B \rightarrow C\}$$

$$- \text{for } f: A \times B \longrightarrow C \text{ define}$$

$$\lambda f: A \longrightarrow (B \Rightarrow C)$$

$$a \longmapsto \lambda b. f(a, b)$$

- EXERCISE check $\oplus$

CASE STUDY

Recursive types can be modelled
by initial algebras

– for details see slides, plus...

Illustrative notes

- Take $D = K_1 (\text{Int} \times \text{Unit})$
 $| K_2 (\text{Int} \times D)$

silly lists

eg
(3, (4, (5, ())))
minus constructors K_i

- $F : \text{Set} \rightarrow \text{Set}$ defined from D

$$F X = (\mathbb{Z} \times \underline{1}) + (\mathbb{Z} \times X)$$

complex definition in the slides...

— Define $I \stackrel{\text{def}}{=} \{E \mid E :: D\}$

— Then $F I \xrightarrow{\sigma} I$ is initial
 definition in the slides

— Roughly speaking

$$(\mathbb{Z} \times 1) + (\mathbb{Z} \times I) \xrightarrow{\sigma} I$$

$$(\underline{z}, *) \mapsto K_1(\underline{z}, ())$$

$$(\underline{z}, E) \mapsto K_2(\underline{z}, E)$$

where $\underline{z} :: \text{Int}$ for each $z \in \mathbb{Z}$

- initiality by example

$$(\mathbb{Z} \times 1) + (\mathbb{Z} \times I) \xrightarrow{\sigma} I$$

$$\begin{array}{ccc}
 (z, E) \mapsto K_2(\underline{z}, E) & & \\
 \downarrow & & \downarrow \\
 (z, *) \mapsto K_1(\underline{z}, ()) & & \\
 \downarrow & & \downarrow \\
 (z, *) \mapsto \underline{z} & & \downarrow \\
 (z, \bar{f}E) \mapsto (z + \bar{f}E) & &
 \end{array}$$

$\bar{f} K_1(\underline{z}, ()) \parallel \underline{z}$
 $\bar{f} K_2(\underline{z}, E) \parallel (z + \bar{f}E)$

$F\bar{f}$ (left vertical arrow)
 f (right vertical arrow)

$$(\mathbb{Z} \times 1) + (\mathbb{Z} \times X) \longrightarrow X$$

$$f = [\pi_{\mathbb{Z}}, +]$$

Take $X \stackrel{\text{def}}{=} \mathbb{Z}$

Hence if $\xi = k_2(\underline{3}, k_2(\underline{4}, k_1(\underline{8}, ())))$
then we have

$$\begin{aligned}\bar{f}(\xi) &= 3 + \bar{f} k_2(\underline{4}, k_1(\underline{8}, ())) \\ &= 3 + 4 + \bar{f} k_1(\underline{8}, ()) \\ &= 3 + 4 + 8 \\ &= 3 + 12 \\ &= 15 \in \mathbb{Z} \quad (= X)\end{aligned}$$

CASE STUDY

Colimits - Building Initial Algebras

- for details see slides...

The end

$$\int_C R(LC, LC)$$